

Spectral lags and the energy dependence of pulse width in gamma-ray bursts: contributions from the relativistic curvature effect

Rong-feng Shen [★], Li-ming Song, and Zhuo Li

Particle Astrophysics Center, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100039, China

Accepted 2005 April 27. Submitted 2004 July 20

ABSTRACT

We compute the temporal profiles of the gamma-ray burst pulse in the four Burst and Transient Source Experiment (BATSE) Large Area Detector (LAD) discriminator energy channels, with the relativistic curvature effect of an expanding fireball being explicitly investigated. Assuming an intrinsic “Band” shape spectrum and an intrinsic energy-independent emission profile, we show that merely the curvature effect can produce detectable spectral lags if the intrinsic pulse profile has a gradually decaying phase. We examine the spectral lag’s dependences on some physical parameters, such as the Lorentz factor Γ , the low-energy spectral index, α , of the intrinsic spectrum, the duration of the intrinsic radiation t'_d and the fireball radius R . It is shown that approximately the lag $\propto \Gamma^{-1}$ and $\propto t'_d$, and a spectrum with a more extruded shape (a larger α) causes a larger lag. We find no dependence of the lag on R . Quantitatively, the lags produced from the curvature effect are marginally close to the observed ones, while larger lags require extreme physical parameter values, e.g., $\Gamma < 50$, or $\alpha > -0.5$. The curvature effect causes an energy-dependent pulse width distribution but the energy dependence of the pulse width we obtained is much weaker than the observed $W \propto E^{-0.4}$ one. This indicates that some intrinsic mechanism(s), other than the curvature effect, dominates the pulse narrowing of GRBs.

Key words: relativity – gamma-rays: bursts – gamma-rays: theory

1 INTRODUCTION

Cheng et al. (1995) were the first to analyse the spectral lag of gamma-ray bursts (GRBs), which is the time delay between the peaks in the Burst and Transient Source Experiment (BATSE) Large Area Detector (LAD) Channel 1 (25 - 50 keV) and Channel 3 (100 - 300 keV) light curves. Subsequently, several authors have carried out more analysis work on the GRB lags. Norris et al. (1996) and Norris, Marani & Bonnell (2000) found the cross-correlation function lags between BATSE Channel 1 and Channel 3 photons tend to concentrate near < 100 ms; for six bursts with known redshift, the z -corrected lags are distributed between

6 and 200 ms. Wu & Fenimore (2000) extended the analysis to very low energy (~ 2 keV); they found that about 20% of GRBs have detectable lags and that GRBs don’t show larger lags at lower energy. A recent measurement by Chen et al. (2005) for the BATSE bursts shows that the majority of the lags are below ~ 200 ms, and that the histogram of the lags peaks around 30 ms. More intriguingly, Norris et al. (2000) found that, for those six bursts with known z , the peak luminosity is anti-correlated with the lags. This relationship provides a useful tool to estimate the distances of large sample of GRBs by analysing their light curves.

Three theoretical explanations for the lag/luminosity relation have been proposed: the relationship is due to the variation in line-of-sight velocity among bursts (Salmonson 2000); it is caused by the variation of the off-axis angle when viewing a narrow jet (Ioka & Nakamura 2001); or it is caused by radiation cooling - highly luminous burst cools fast and

[★] E-mail: rfshen@astro.as.utexas.edu. Present address: Department of Astronomy, University of Texas at Austin, Austin, TX 78712, USA

the lag will be short (Schaefer 2004). However, the problem of what mechanism(s) causes the spectral lags of GRBs remains unresolved. Salmonson (2000) did not explain the origin of the lag but assumed that it derives from some proper decay time scale $\Delta t'$ in the rest frame of the emitter. In the model by Ioka & Nakamura (2001), the lag is caused by the far side of the emitting region producing lower-energy radiation after a longer light-travel time, for a narrow jet with viewing angles outside the cone of jet. However, their model requirements seem too stringent (see below). Schaefer (2004) proposed radiative cooling as the origin of the lag. The difficulty of this explanation is that in order to adjust the observed synchrotron cooling time scale to be comparable with the lag time scale, the magnetic field has to be ~ 7 Gauss, a value much below the strength required by most of the current models (e.g. Piran 1999).

Kocevski & Liang (2003) have assumed that the observed lag is the direct result of spectral evolution, another property of GRBs (Norris et al. 1986; Bhat et al. 1994). In particular, as the peak energy of the GRB's νF_ν spectrum decays through the four BATSE channels, the photon-flux peak in each individual channel will be shifted, probably producing the measured lag. From a sample of 19 GRBs, Kocevski & Liang (2003) found an empirical relation between the peak energy's decaying rate and the GRB lag.

It is widely accepted that the gamma-rays come from a relativistically expanding fireball surface with Lorentz factor $\Gamma > 100$ (Lithwick & Sari (2001) and references therein). At some distance from the central source (e.g. $R = 10^{12} \sim 10^{14}$ cm, *cf.* Piran (1999)), photons emitted from the region on the line of sight and those from the side region at an angle of $\theta \sim 1/\Gamma$ with respect to the line of sight are Doppler-boosted by different factors and travel different distances to the observer. This is what we call the curvature effect. Comparing the radiation from the side region and that from the line-of-sight region, for the latter its photons are Doppler-boosted to higher energies and arrive to the observer earlier; its observed temporal structure will be boosted to be narrower.

The motivation of this paper is to see how the curvature effect will change the intrinsic pulse profile in different energy channels, and special interest is focused on whether merely the curvature effect can produce the spectral lags of the pulses. Except for the soft photon lags, the pulses in GRBs show another temporal property, i.e. the pulse narrowing at higher energy, or pulse width as a function of energy (Fenimore et al. 1995, Norris et al. 1996). We are also interested in probing the contributions of the curvature effect to these properties.

A fireball internal-external shocks model has emerged for the theoretical understanding of the origin of GRBs (Piran 1999, 2004). According to this model, GRBs are produced when an ultra-relativistic outflow dissipates its kinetic energy through the internal collisions within the outflow itself. The afterglow occurs when the flow is decelerated by shocks with the circumburst medium. This model has made many successful explanations and predictions to the obser-

vations of GRBs. Our investigation to the curvature effect will be based on the frame of the internal shock model, where the shock is generated from the colliding shells, and electrons are accelerated in the shock and radiate.

Ioka & Nakamura (2001) proposed a model in which a narrow jet is viewed at off-axis angle to explain the lag/luminosity relation and the variability/luminosity relation. Their model successfully reproduces the lag/luminosity relation, while the lag is caused by the curvature effect of the jet, which increases with the off-axis angle. However, one substantial problem with this model, as pointed out by Schaefer (2004), is that it works with an exacting assumption that the jet opening angle always equals Γ^{-1} . Furthermore, Ioka & Nakamura (2001) only consider an instantaneous emission in the jet rest frame, which is a too simple assumption. Consequently, in their model there is no lag when the jet is viewed near-on-axis. Different from Ioka & Nakamura (2001), we have considered various rest-frame emission profiles and assumed an isotropically expanding radiation surface.

We present our model in Section 2, including the basic assumptions and formulas. The major results are presented in Section 3, based on which we give our conclusions and discussion in Section 4.

2 MODEL

2.1 Three time scales

Three time scales are involved in determining the temporal structures of pulses in GRBs: (i) cooling time scale; (ii) hydrodynamic time scale; (iii) angular spreading time scale (Kobayashi et al. 1997, Wu & Fenimore 2000).

In the synchrotron cooling model, the shock-accelerated electrons cool via synchrotron emission, and the electron's average energy becomes smaller and the radiated power decays. As pointed out by Wu & Fenimore (2000), the standard internal-shock model gives an observed synchrotron cooling time scale at a given photon energy as

$$T_{syn}(h\nu) \sim 2 \times 10^{-6} s \epsilon_B^{-3/4} \left(\frac{h\nu_{obs}}{100 keV} \right)^{-1/2},$$

where ϵ_B is the equipartition parameter for the ratio of the magnetic energy density to the total internal energy density; for a typical value in the internal-shock model, e.g. $\epsilon_B = 0.01$ ($B \sim 10^5$ G), the cooling time scale is far shorter than the lag time scale.

The hydrodynamic time scale is related to the energizing of the electrons. In the internal-shock model, if one assume the the local microscopic acceleration of electrons is instantaneous, then the hydrodynamic time scale is attributed to the shell-crossing time of the shock, $T'_{dyn} = \Delta'/v'_{sh}$, where Δ' is the shell width and v'_{sh} is the shock velocity, both in the comoving frame of the upstream flow. We hardly know about Δ' . However if one assume the shells are radially ex-

panding, in the observer frame this time scale is

$$T_{dyn} \sim 1s \beta_{sh}'^{-1} \left(\frac{R}{10^{15} cm} \right) \left(\frac{\Gamma}{100} \right)^{-2},$$

where R is the radius at which the shell radiates, and Γ is the Lorentz factor of the shell (Ryde & Petrosian 2002). (Note that apart from the shock acceleration scenario, there can be other particle energizing mechanisms, e.g. magnetic field reconnection (Stern 1999), which may have a different hydrodynamic time scale)

The angular spreading time scale is the delay between the arrival times of the photons emitted at the line-of-sight region and of that emitted at the side region of the shell (Sari & Piran 1997). Because of the relativistic beaming of the moving radiating particles, only the emission from a narrow cone with an opening angle of $\sim 1/\Gamma$ is observed. This gives a time scale of the delay of

$$T_{ang} = 1.7s \left(\frac{R}{10^{15} cm} \right) \left(\frac{\Gamma}{100} \right)^{-2}.$$

2.2 Assumptions

In this paper, different from Schaefer (2004), we assume the cooling time scale is much shorter than the other two time scales, i.e. the accelerated particles radiate their energy rapidly. Thus the rest frame duration of the emission is determined by the hydrodynamic time scale.

We consider a thin shell expanding with a relativistic speed, whose Lorentz factor is Γ . The shell begins to radiate at radius R . In the co-moving frame of the shell, the radiation intensity of the shell surface $I'(\nu', t')$ is assumed to be isotropic, and has an energy-independent time history $f'(t')$. Note that all the quantities in the rest frame of the radiation surface are labeled with a prime note.

Band et al. (1993) found that the GRB spectra are well described at low energy by a power law with an exponential cutoff and by a steeper power law at high energy. The typical fitted value distributions for the low-energy spectral index (α) is $-1.5 \sim -0.3$, the high-energy spectral index (β) is $-2 \sim -3$ and the peak energy (E_p) of the νF_ν spectrum is $100 \sim 500$ keV (see Preece et al. 2000). By modeling, Qin (2002) showed that the relativistic expanding of the fireball does not alter the shape of GRB's rest frame spectrum, but only shifts the peak of the spectrum to a higher energy; the distribution of E_p is scaled with the Lorentz factor Γ , as shown in their Table 4. So we directly adopt the spectral form used by Band et al. (1993) as the rest frame spectrum, and choose values for the low-energy spectral index α to be -0.8 and the high-energy spectral index β be -2.4 ; The peak energy of the rest frame spectrum E_p' is adjusted such that E_p is 350 keV, where the relation $E_p = 1.65 \times \Gamma E_p'$, derived from Table 4 in Qin (2002) in the range of $10 < \Gamma < 2000$, is used.

2.3 The formulas

The radiation intensity in the observer's frame is $I(\mu, \nu, t) = I(\mu, \nu) f(t)$. It is connected with the rest frame radiation intensity by

$$I(\mu, \nu) f(t) = \mathcal{D}^3(\mu) I'(\nu') f'(\Gamma t') \quad (1)$$

and

$$\nu = \nu' \mathcal{D}(\mu), \quad (2)$$

where $\mu = \cos\theta$, and θ is the angle of the concerned local radiation surface to the line of sight; $\mathcal{D}(\mu) = [\Gamma(1 - \mu\beta)]^{-1}$ is the local relativistic Doppler factor respect to the observer, where Γ is the Lorentz factor of the radiation surface and $\beta = \sqrt{1 - \Gamma^{-2}}$.

We use t to refer to the photon emitting time and use T to refer to the time that the photon arrives at the observer. We define that, at the time $t = 0$, the first photons are emitted from the surface and, for simplicity, we also tune the arrival time T of the first photon emitted from the line-of-sight region ($\mu = 1$) to be 0 too. Then the arrival time of a photon emitted from region μ at time t is

$$T = (1 - \beta)t + (1 - \mu)(R + \beta ct)/c.$$

The first part of the right-hand side of the equation is caused by the motion of the shell, and the second part is due to the difference between the light travel distances of photons emitted from the line-of-sight region and from the side region. It can be rewritten as

$$T = (1 - \mu\beta)t + (1 - \mu)\tau, \quad (3)$$

where $\tau = R/c$, which connects the photon's arrival time, emitting time and emitting place in one equation.

At time T , the observed flux comes from the photons emitted at the region $1 > \mu > \mu(T, t = 0)$, where the boundary $\mu(T, t = 0) = 1 - T/\tau$, calculated from equation (3), is the place whose first photon is emitted with arrival time T . The observed specific flux can be obtained by integrating the radiation intensity over this region

$$F_\nu(T) = - \int_{1-T/\tau}^{1-T/\tau} I_\nu(\mu, t) \mu R^2(t) d\mu \quad (4)$$

For a BATSE LAD discriminator channel (ν_a, ν_b), the observed photon counts flux is

$$n_{ab}(T) = \int_{\nu_a}^{\nu_b} \frac{F_\nu(T)}{\nu} d\nu.$$

From equation (4), substituting the emitting time t in the integral with T and μ using equation (3), $n_{ab}(T)$ can be rewritten as

$$n_{ab}(T) = -[\tau + (T - \tau)\beta]^2 \int_{1-T/\tau}^{1-T/\tau} \times \frac{f[t(T, \mu)]}{(1 - \mu\beta)^2} \mu d\mu \mathcal{D}^3(\mu) \int_{\nu_a/\mathcal{D}(\mu)}^{\nu_b/\mathcal{D}(\mu)} \frac{I'_{\nu'}}{\nu'} d\nu',$$

where $R(t) = c[\tau + t(T, \mu)\beta]$ and equations (1), (2) are used. So we get the explicit formula that is used in our calculation.

2.4 Intrinsic time profile of the radiation

Here we introduce three cases of the intrinsic emission time profile.

(i) Rectangular profile. During a finite duration, the emissivity is constant, with the instantaneous rising phase and decaying phase:

$$f(t) = \begin{cases} 1, & 0 < t < t_d \\ 0, & t > t_d \end{cases}$$

In our assumptions t_d is related to the hydrodynamic time scale discussed above, but note that t_d is defined in the observer's frame, without taking into account the Doppler boosting (i.e., it is not the observed time scale). In addition, we refer to t'_d as the t_d measured in the comoving frame of the shell.

(ii) One-sided exponential profile. The emissivity decays exponentially after an instantaneous rising phase:

$$f(t) = \begin{cases} 0, & t < 0 \\ \exp[-(t/t_d)], & t > 0 \end{cases}$$

(iii) Symmetric Gaussian profile. The emissivity has both a finite-time rising phase and a finite-time decaying phase:

$$f(t) = \begin{cases} 0, & t < 0 \\ \exp[-(\frac{t-1.5t_d}{t_d})^2], & t > 0 \end{cases}$$

where we introduce the coefficient -1.5 in order that the emission starts at $t = 0$ with the radiation intensity of a tenth ($e^{-1.5} \approx 0.1$) of its peak value.

3 RESULTS

3.1 Observed temporal profiles in different energy bands and time lags

Using the equations and the typical parameters introduced above, we calculate the observed pulse profiles in the four BATSE LAD energy channels, assuming three different intrinsic pulse profiles. The results are plotted in Figure 1, 2 and 3, respectively.

First, we calculate the pulse light curves in the four energy channels for the rectangular intrinsic radiation profile. It has a steady rising phase, followed by a distinct peak, as shown in Figure 1. The rising of the flux is due to the expanding of the radiation surface. The rising is steady because the radiation intensity is constant during a finite time. The decay phase of the observed pulse is due to the angular spreading effect. The peak occurs when the intrinsic radiation begins to cease. For various parameter spaces (duration of the emission $t_d : 10^3 - 10^5$ s, radius $R : 10^{13} - 10^{15}$ cm, Lorentz factor $\Gamma : 50 - 500$), we do not detect any time lag between the peaks observed at different energies in this case.

As for the one-sided exponential and the symmetric Gaussian emission profiles, their observed light curves calculated in the four energy channels are illustrated in Figure




Figure 1. Observed pulse from intrinsic rectangular time profile in four BATSE LAD energy channels. *Solid line:* 25 – 50 keV; *dashed line:* 50 – 100 keV; *dot-dashed line:* 100 – 300 keV; *dotted line:* 300 – 1800 keV. The vertical coordinate is in arbitrary units. The peaks of the pulses at 4 energies arrived simultaneously. $\Gamma = 100$, $\alpha = -0.8$, $\beta = -2.4$, $R = 3 \times 10^{14}$ cm, $t_d = 0.9 \times 10^4$ s. The rising phase comes from the expanding of the radiation surface and the rising time is determined by duration of the intrinsic radiation. The decay phase is due to the angular spreading effect.

2 and 3, respectively. For these two emission profiles, their peaks are more gradual. The maximum of the photon flux at higher energy arrives early than at lower energy. If we define the time lag as being the difference between the arrival times of the peaks in energy Channel 1 and 3 or 4, the reproduced lags are quantitatively comparable to the observed ones (e.g., $\sim 10^{-2}$ s if corrected for the cosmological time dilation. cf. Norris et al. (2000)). Note that here we do not take into account the redshifts, z , of the GRBs, which must make the observed lags $(1+z)$ times larger.

The intrinsic rectangular profile cannot produce the observed lags because in this case the radiation diminishes immediately. In the observed pulse, the rising phase comes from the expanding of the radiation surface and the rising time is determined by duration of the intrinsic radiation ($\approx t_d/(2\Gamma^2)$; also see Qin et al. 2004). The decay phase is due to the angular spreading effect. Hence if the intrinsic radiation switches off immediately, the transition from the rising phase to the decay occurs abruptly and induces a sharp peak in the observed pulse. For this case, the immediate switch-off of the intrinsic radiation dilutes the relativistic curvature effect, and hence does not produce the peak lags.

3.2 The pulse width - energy relation

For the three intrinsic emission profiles, we find that the pulse observed in high-energy channel is narrower than in lower-energy channel, which is manifested by the fact that the FWHMs of the pulses in separated energy channels decrease with the energy in a power-law form. However, the pulse narrowing we obtained is less prominent than that ob-




Figure 2. Observed pulse from one-sided exponential intrinsic time profile in four BATSE LAD energy channels. *Solid line*: 25 – 50 keV; *dashed line*: 50 – 100 keV; *dot-dashed line*: 100 – 300 keV; *dotted line*: 300 – 1800 keV; *triple-dots-dashed line*: the intrinsic emission profile. The vertical coordinate is in arbitrary units. $\Gamma = 400$, $\alpha = -0.8$, $\beta = -2.4$, $R = 5 \times 10^{14}$ cm, $t_d = 5 \times 10^4$ s. $\text{Lag}_{13} \simeq 0.03$ s.




Figure 3. Observed pulse from symmetric Gaussian intrinsic time profile in four BATSE LAD energy channels. *Solid line*: 25 – 50 keV; *dashed line*: 50 – 100 keV; *dot-dashed line*: 100 – 300 keV; *dotted line*: 300 – 1800 keV; *triple-dots-dashed line*: the intrinsic emission profile. The vertical coordinate is in arbitrary units. $\Gamma = 400$, $\alpha = -0.8$, $\beta = -2.4$, $R = 5 \times 10^{14}$ cm, $t_d = 2 \times 10^4$ s. $\text{Lag}_{13} \simeq 0.005$ s.

served in real GRBs for which pulse width decay power-law index ~ -0.4 ; while the pulse width decay index we obtained, for instance in the case of Figure 2 (one-sided exponential decay emission profile), is -0.13 .

Other than the “Band” spectrum, we also used an alternative function - a low-energy power law plus the high-energy exponential cut-off at E'_p - for the rest-frame spectrum. For the typical values of the parameters we have used, this changing of spectrum only narrows the Channel 4 pulse




Figure 4. Symmetric Gaussian intrinsic pulse: lag dependence on the Lorentz factor. $\alpha = -0.8$, $\beta = -2.4$, $R = 5 \times 10^{13}$ cm, $t'_d = 20$ s. Power-law fit gives index -0.87 and -0.93 , respectively.

by $\sim 6\%$, hence hardly changes the slope of the pulse width versus the energy.

3.3 Lag's dependence on other physical parameters

The spectral lag is an important observational property of the pulse in GRBs in that it may be used to derive the cosmological distribution of GRBs (Norris 2002) and to discriminate the internal shock signature and the external shock signature in the pulses (e.g., Hakkila & Giblin 2004). This motivates us to probe the dependences of the peak lags on other physical parameters of the simple model. We choose the symmetric Gaussian profile as the intrinsic emission profile, which includes an intrinsic rising phase. We alter the Lorentz factor Γ , the spectral parameters of the rest frame emission (α , β and E'_p), the radius of the radiation surface R , and the rest-frame duration t'_d of the intrinsic radiation, respectively, and see how the Channel 1/3 and the Channel 1/4 peak lags vary with these changes.

3.3.1 Lorentz factor

Figure 4 shows that the lag decreases with the Lorentz factor following $\text{Lag} \propto \Gamma^{-1}$. We think this is a natural outcome of the relativistic boosting of the time structure. Those pulses whose lags are larger may come from colliding shells with low Lorentz factors, according to the current standard models (e.g., Piran 1999).

3.3.2 Spectral parameters

Figure 5 shows that the lag increases with the low energy spectral index α of the rest frame spectrum of the pulse. A larger α means more photons are concentrated around the peak energy E'_p of the $\nu' F'_\nu$ spectrum. Then, for a narrower




Figure 5. Symmetric Gaussian intrinsic pulse: lag dependence on α , the low-energy spectral index of the adopted spectrum. $\Gamma = 100$, $E'_p = 1.75$ keV, $\beta = -2.4$, $R = 5 \times 10^{13}$ cm, $t'_d = 40$ s. Note that GRB spectrum with $\alpha > 0$ is very rarely observed (Preece et al. 2000); the calculated data points there are only used to show the tendency.




Figure 6. Symmetric Gaussian intrinsic pulse: lag dependence on the observed spectral break energy E_p of the pulse spectrum. $\Gamma = 400$, $\alpha = -0.8$, $\beta = -2.4$, $R = 5 \times 10^4$ cm, $t'_d = 100$ s. Note that E_p is obtained through a simple scaling relation, $E_p = 1.65 \times \Gamma E'_p$, between E_p and the break energy, E'_p , of the rest-frame emission spectrum, as a result of the simulation by Qin (2002).

spectrum, the curvature effect will work more effectively in producing the spectral lags. This conjecture is supported when we alter the high-energy spectral index β . We found that a steeper high-energy power-law spectrum produces a larger lag, e.g., the Channel 1/4 lag has a 14% increase for β changing from -2.4 to -3.0. Compared with the Channel 1/4 lag, the Channel 1/3 lag has a weaker dependence upon β .

The lag's dependence on the observed break energy of the spectrum E_p is shown in Figure 6. The lag has its maximum when E_p falls near the starting energy of the corresponding high-energy channel that is used in measuring the

lag (i.e., ~ 100 keV for Channel 3, ~ 300 keV for Channel 4).

The above findings about the lag's dependences on the spectral parameters are qualitatively consistent with the tendency observed in those long-lag wider-pulse bursts by Norris et al. (2005). They found that their long-lag (measured for Channel 1/3) burst sample has, on average, lower E_p (centered around ~ 110 KeV), larger α (harder low-energy power law) and smaller β (softer high-energy power law), than the bright burst sample analyzed by Preece et al. (2000).

Substituting the “Band” spectrum with an alternative one of a single power law plus an exponential high-energy cut-off causes no changes to the Channel 1/3 lag, while the Channel 1/4 lag has a $\sim 40\%$ increase if the observed cut-off energy E_p is below 300 keV; for $E_p > 300$ keV, the increase of Channel 1/4 lag is much smaller.

3.3.3 Duration of the emission

We find that longer rest-frame duration (t'_d) of emission will cause larger lags, as is shown in Figure 7. This result may be associated with an observed tendency that wider pulses exhibit longer lags (Norris et al. 1996; Norris, Scargle & Bonnell 2001; Norris et al. 2005).

3.3.4 Radius of the radiation surface

The lag appears to be independent on R , the radius of the radiation surface (see Figure 8). We know R determines the angular spreading time scale, $T_{ang} \approx R/(2\Gamma^2 c)$. This result suggests that though in our model the angular spreading effect is a necessity in causing the lags, the peak lag in the pulses is not correlated with the angular spreading time scales.

In addition, we find that the decrease of pulse width with the photon energy is dependent upon the low-energy spectral index of the radiation spectrum, as shown in Figure 9.

4 CONCLUSIONS AND DISCUSSIONS

By assuming an intrinsic “Band”-shape spectrum and an exponential or Gaussian emission profile, we show that merely the curvature effect produces detectable soft lags in the GRB pulses. Therefore the soft time lags can be a signature of the relativistic motion occurring in GRBs.

The observed Channel 1/3 lags are typically distributed among $10^{-2} - 10^{-1}$ s (Norris et al. 2000). For typical physical parameters, i.e., $\Gamma \approx 100$, $t'_d \approx 40$ s, $\alpha \approx -1$, as shown in Figure 4 - 8, the lags produced by the relativistic curvature effect are slightly above 10^{-2} s, marginally close to those observed, after considering the cosmological time dilation effect if a GRB redshift of 2 is assumed. To account for those observed larger lags (~ 0.1 s), it requires extreme physical parameter values, e.g., $\Gamma < 50$ or $\alpha > -0.5$.




Figure 7. Symmetric Gaussian intrinsic pulse: lag dependence on t'_d , the rest frame duration of the radiation. $\Gamma = 100$, $E'_p = 1.75$ keV, $\alpha = -0.8$, $\beta = -2.4$, $R = 5 \times 10^{13}$ cm.




Figure 8. Symmetric Gaussian intrinsic pulse: lag dependence on R , the radius of radiation surface. $\Gamma = 100$, $E'_p = 1.75$ keV, $\alpha = -0.8$, $\beta = -2.4$, $t'_d = 40$ s.

We did not find any peak lag for a rectangular intrinsic emission profile, from which a straight-forward conclusion regarding the radiation process of the pulse in GRBs can be obtained — the radiation intensity must have a decaying phase in order to produce the observed peak lags. The intrinsic decaying phase may be due to the variations associated with hydrodynamic processes, such as the decaying of emission caused by density or magnetic field attenuation as the shock moves through the shell.

We have investigated the possible dependences of the pulse peak lag upon other physical parameters of the kinematic model. We found: the lag is proportional to the inverse of the Lorentz factor; the lag is proportional to the duration




Figure 9. Symmetric Gaussian intrinsic pulse: pulse width as a function of the photon energy for different α , the low energy spectral index of the adopted spectrum. Other model parameters are the same as in Figure 5.

of the intrinsic radiation t'_d ; the lag is weakly dependent on R ; the lag is larger when larger amount of energy is concentrated at E_p (larger α or smaller β).

The pulse width decreases with energy ($W \propto E^{-0.1 \sim -0.2}$), but not as fast as the observed ($W \propto E^{-0.4}$), though we found a faster decrease with a larger low-energy spectral index α . There must be other energy-dependent narrowing mechanisms underlying. Similar to this conclusion, Dermer (2004) pointed out that other processes including adiabatic and radiative cooling, a non-uniform jet, or the external shock process, rather than the curvature effect, should be needed to explain the relationship between the measured peak photon energy E_p and the measured νF_ν flux at E_p in the decaying phase of a GRB pulse. They found the curvature relationship does not agree with the observation (Borgonovo & Ryde 2001).

For simplicity we have assumed a spherically symmetric radiation surface in this paper, though there is some observational evidence indicating that the GRB outflow may be collimated. Derived from the afterglow observations, the jet coming from the GRB central source generally has a half opening angle $\theta_j > \Gamma^{-1}$ (Frail et al. 2001). Assuming a jet geometry with the jet opening angle of 1° or 4° and the same parameter values (Γ , R and t'_d) used in the spherical geometry, we calculated the observed pulse shapes and the lags; they show no difference from those of the spherical geometry. The reason for this is as follows. Even in the case of the isotropic radiation surface, the contribution of the flux from the outer side region where the observing angle θ is larger than Γ^{-1} is relatively very small, because the local flux contribution from the radiation surface (i.e., the first-part integrand of Equation 5 in Section 2) will decrease drastically with θ as $\propto (1 + \Gamma^2 \theta^2)^{-5}$ when θ is small.

ACKNOWLEDGMENTS

We thank the referee for their careful reading of the manuscript and for many valuable suggestions. This work was supported in part by the Special Funds for Major State Basic Science Research Projects of the Ministry of Science and Technology, and by the National Natural Science Foundation of China through grant 10473010.

REFERENCES

- Band D., Matteson J., Ford L. et al. 1993, *ApJ*, 413, 281
 Bhat P. N., Fishman G. J., Meegan, C. A., et al. 1994, *ApJ*, 426, 604
 Borgonovo L. & Ryde F. 2001, *ApJ*, 548, 770
 Chen L., Lou Y.-Q., Wu M., et al. 2005, *ApJ*, 619, 983
 Cheng L. X., Ma Y. Q., Cheng K. S., Lu T. & Zhou Y. Y. 1995, *A&A*, 300, 746
 Dermer C. D. 2004, *ApJ*, 614, 284
 Fenimore E. E., in t Zand J. J. M., Norris J. P., et al. 1995, *ApJ*, 448, L101
 Frail D. A., Kulkarni S. R., Sari R. et al. 2001, *ApJ*, 562, L55
 Hakkila J. & Giblin T. W. 2004, *ApJ*, 610, 361
 Ioka K. & Nakamura T. 2001, *ApJ*, 554, L163
 Kobayashi S., Piran T. & Sari R. 1997, *ApJ*, 490, 92
 Kocevski D. & Liang E. 2003, *ApJ*, 594, 385
 Lithwick Y. & Sari R. 2001, *ApJ*, 555, 540
 Norris, J. P., Share, G. H., Messina, D. C. et al. 1986, *ApJ*, 301, 213
 Norris J. P., Nemiroff R. J., Bonnell J. T. et al. 1996, *ApJ*, 459, 393
 Norris J. P., Marani G. F. & Bonnell J. T. 2000, *ApJ*, 534, 248
 Norris J. P., Scargle, J. D. & Bonnell, J. T., in Ritz S., Gehrels N., Shrader C. R., eds, *AIP Conf. Proc. Vol. 587, Gamma 2001: Gamma-Ray Astrophysics*. American Institute of Physics, New York, p. 176 (astro-ph/0105052)
 Norris J. P. 2002, *ApJ*, 579, 386
 Norris J. P., Bonnell J. T., Kazanas D. et al. 2005, *ApJ*, 627, 324
 Piran T. 1999, *Phys. Rep.*, 314, 575
 Piran T. 2004, *Rev. Mod. Phys.*, 76, 1143
 Preece R. D., Briggs M.S., Mallozzi R. S., et al. 2000, *ApJS*, 126, 19
 Qin Y.-P. 2002, *A&A*, 396, 705
 Qin Y.-P., Zhang Z.-B., Zhang F.-W., et al. 2004, *ApJ*, 617, 439
 Ryde F. & Petrosian V. 2002, *ApJ*, 578, 290
 Salmonson J. D. 2000, *ApJ*, 544, L115
 Sari R. & Piran T. 1997, *ApJ*, 485, 270
 Schaefer B. 2004, *ApJ*, 602, 306
 Stern B. in: Poutanen J, Svensson R, eds. *High Energy Processes in Accreting Black Holes*, ASP Conference Series 161. San Francisco: ASP, 1999, p. 277
 Wu B. & Fenimore E. 2000, *ApJ*, 535, L29